

University of Groningen

On boson condensation into an infinite number of low-lying levels

van den Berg, M

Published in:
Journal of Mathematical Physics

DOI:
[10.1063/1.525445](https://doi.org/10.1063/1.525445)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1982

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van den Berg, M. (1982). On boson condensation into an infinite number of low-lying levels. *Journal of Mathematical Physics*, 23(6), 1159-1161. <https://doi.org/10.1063/1.525445>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

On boson condensation into an infinite number of low-lying levels

M. van den Berg

Citation: [Journal of Mathematical Physics](#) **23**, 1159 (1982); doi: 10.1063/1.525445

View online: <https://doi.org/10.1063/1.525445>

View Table of Contents: <http://aip.scitation.org/toc/jmp/23/6>

Published by the [American Institute of Physics](#)

PHYSICS TODAY

WHITEPAPERS

MANAGER'S GUIDE

Accelerate R&D with
Multiphysics Simulation

READ NOW

PRESENTED BY

 **COMSOL**

On boson condensation into an infinite number of low-lying levels

M. van den Berg

Institute for Theoretical Physics, State University of Groningen, P. O. Box 800, Groningen, The Netherlands

(Received 20 November 1981; accepted for publication 25 January 1982)

The complicated structure of the condensate of a free boson gas in two dimensions with mixed boundary conditions is examined.

PACS numbers: 05.30.Jp

I. INTRODUCTION AND THEOREM

It has been shown in recent papers^{1,2} that there exist three types of condensation in the free boson gas:

- (I) Macroscopic occupation of a finite number of single-particle levels.
- (II) Macroscopic occupation of an infinite number of single-particle levels.
- (III) Nonextensive condensation (no levels macroscopically occupied).

In this paper we will present an example of type II. Let B_L be a finite region in R^d with volume L^d and let $E_1 < E_2^L < E_3^L \dots$ be the spectrum of the single-particle Hamiltonian H_L ; then [if we assume that $\exp(-\beta H_L)$ is trace-class] the mean number $\langle n_k \rangle_L$ of particles per volume in the k th state for a free boson gas is given by (in the grand canonical ensemble³)

$$\langle n_k \rangle_L = \frac{1}{L^d} \frac{\zeta(L)}{e^{\eta_k^L} - \zeta(L)}, \quad (1)$$

where $\eta_k^L = E_k^L - E_1^L$ and $\zeta(L)$ is the positive solution of

$$\sum_k \langle n_k \rangle_L = \rho; \quad (2)$$

ρ is the mean number of particles per unit volume. The thermodynamic limit is the limit in which $L \rightarrow \infty$ and the number density ρ is kept fixed. If the spectrum of H_L is such that the critical density ρ_c , defined by

$$\rho_c = \lim_{\zeta \uparrow 1} \lim_{L \rightarrow \infty} \frac{1}{L^d} \sum_k (e^{\eta_k^L} - \zeta)^{-1}, \quad (3)$$

is finite, then condensation of type I, II, or III takes place if $\rho > \rho_c$. In the case when $H_L = -\Delta/2$ (with Dirichlet or Neuman conditions on the boundary ∂B_L of B_L) the critical density ρ_c is finite only for $d > 2$. However, if we impose attractive boundary on the eigenfunctions, then ρ_c is finite for all dimensions $d = 1, 2, \dots$. This was noticed by Robinson⁴ and discussed in detail by Landau and Wilde.⁵ The reason that ρ_c is finite in this case is that the attractive boundary conditions cause a gap in the single-particle spectrum:

$$E_j^L - E_1^L > g, \quad \text{for some } j > 1 \text{ and all } L > L_0. \quad (4)$$

It was noticed before by Bijl, De Boer, and Michels⁶ that a gap in the spectrum will change the thermodynamical properties.

In this paper we will take $H_L = -\Delta/2$ and impose position-dependent boundary conditions. We will see that they lead to condensation of type II for $d = 2$ and of type I for $d = 1, d = 3, \dots$. If the boundary conditions are position-independent, then there is condensation for $\rho > \rho_c$ of type I

only (as in Ref. 4). Let us take for B_L the cuboid

$$B_L = \{x \in R^d: 0 < x_1 < L, \dots, 0 < x_d < L\} \quad (5)$$

and let the boundary conditions be

$$\frac{\partial \phi}{\partial x_1} = -\sigma \phi \quad \text{for } x_1 = L, 0 < x_2 < L, \dots, 0 < x_d < L, \quad (6)$$

$\phi = 0$ for any other point of the boundary.

The boundary conditions are said to be attractive if $\sigma < 0$. We have expressed all lengths in units $(\beta \hbar^2/m)^{1/2}$, where β is the inverse of the temperature times Boltzmann's constant. Let $\{\Phi_k^L\}$ be the eigenfunction of the Laplacian in B_L with boundary conditions (6) so that

$$-\frac{1}{2} \Delta \Phi_k^L = E_k^L \Phi_k^L. \quad (7)$$

Then

$$\begin{aligned} \Phi_k^L &= \left(\frac{2}{L}\right)^{d/2} \left(1 - \frac{\sin 2L(2\epsilon_{k_1})^{1/2}}{2L(2\epsilon_{k_1})^{1/2}}\right)^{-1/2} \\ &\quad \times \sin x_1(2\epsilon_{k_1})^{1/2} \prod_{i=2}^d \sin \frac{\pi k_i x_i}{L}, \\ E_k^L &= \epsilon_{k_1} + \sum_{i=2}^d \frac{\pi^2 k_i^2}{2L^2} \\ k &= (k_1, \dots, k_d), \\ k_i &= 1, 2, 3, \dots \quad (i = 1, \dots, d), \end{aligned} \quad (8)$$

and $\{\epsilon_{k_1}\}$ are the positive solutions for ϵ of

$$\tan L(2\epsilon)^{1/2} = -(2\epsilon)^{1/2}/\sigma, \quad (9)$$

ordered so that $\epsilon_1 < \epsilon_2 < \epsilon_3 \dots$. For $\sigma < -1/L$ and $L \gg 1$, the spectrum is approximately given by

$$\epsilon_{k_1} \sim (\pi^2/2L^2)(k_1 - 1)^2, \quad k_1 = 2, 3, \dots, \quad (10)$$

while there is a bound state with energy

$$\epsilon_1 \sim -\sigma^2/2. \quad (11)$$

Our main result is the following:

Theorem: For $\rho < \rho_c$, $\zeta(L)$ tends to the positive solution of

$$\rho = \sum_{n=1}^{\infty} \frac{\zeta^n}{(2\pi n)^{d/2}} e^{-n\sigma^2/2}. \quad (12)$$

For $\rho > \rho_c$, $\zeta(L)$ tends to 1 as follows:

$$\zeta(L) \sim 1 - 1/AL^2, \quad d = 2,$$

$$\zeta(L) \sim 1 - 1/(\rho - \rho_c)L^d, \quad d = 1, 3, \dots,$$

$$\rho_c = \sum_{n=1}^{\infty} \frac{e^{-n\sigma^2/2}}{(2\pi n)^{d/2}}, \quad (13)$$

and A is the unique solution of

$$\sum_{k_2=1}^{\infty} \left(\frac{\pi^2}{2} (k_2^2 - 1) + \frac{1}{A} \right)^{-1} = \rho - \rho_c. \quad (14)$$

We see that due to (13) and (1)

$$\lim_{L \rightarrow \infty} \langle n_k \rangle_L = \begin{cases} [(\pi^2/2) (k_2^2 - 1) + 1/A]^{-1} & k_1 = 1, k_2 = 1, 2, 3, \dots \\ 0, & k_1 = 2, 3, \dots, k_2 = 1, 2, \dots \end{cases} \quad d = 2, \quad (15)$$

$$\lim_{L \rightarrow \infty} \langle n_k \rangle_L = \begin{cases} \rho - \rho_c, & k = (1, \dots, 1) \\ 0, & k \neq (1, \dots, 1) \end{cases} \quad d \neq 2.$$

It follows that in two dimensions there is macroscopic occupation of an infinite set of low-lying single-particle states (condensation of type II) whereas in 1, 3, 4, ... dimensions only the ground state is macroscopically occupied (type I).

This can also be seen by looking at the scaled spatial particle density ν_L defined by

$$\nu_L(u) = \sum_k \frac{\xi(L)}{e^{\eta_k^L} - \xi(L)} [\Phi_k^L(Lu)]^2. \quad (16)$$

One finds that

$$\lim_{L \rightarrow \infty} \nu_L(u) = \begin{cases} (\rho - \rho_c) \delta(1 - u_1) + \rho_c, & d = 1, \rho > \rho_c, \\ \delta(1 - u_1) \sum_{j=1}^{\infty} \frac{2 \sin^2 \pi u_2}{(j^2 - 1) \pi^2/2 + 1/A} + \rho_c, & d = 2, \rho > \rho_c, \\ (\rho - \rho_c) \delta(1 - u_1) \prod_{m=2}^d 2 \sin^2 \pi u_m + \rho_c, & d \geq 3, \rho > \rho_c, \\ \rho, & d \geq 1, \rho < \rho_c, \end{cases} \quad (17)$$

where $u \in B_1$ and $\delta(1 - u_1)$ is the Dirac delta function supported on the hyperplane $u_1 = 1$. We see that the expressions [apart from (12) and (13)] are identical to the corresponding expressions for the free boson gas in the presence of an external field of power form in one direction (see Ref. 1). This is for the following reason: The attractive boundary condition causes a gap in the spectrum and forces the wavefunction to have a maximum near the attractive boundary. The same is true for one-particle Hamiltonian with an external potential with an absolute minimum at the boundary (and Dirichlet boundary conditions). We will give a sketch of the proof using the ideas of Lewis and Pulè.⁷

II. SKETCH OF THE PROOF

Let $f_L(z)$ be defined by

$$f_L(z) = \frac{1}{L^d} \sum_{n=1}^{\infty} z^n \sum_{\{k: k_1=1\}} e^{-n \eta_k^L}. \quad (18)$$

The first step is to show that for $z \in [0, 1]$ we have

$$\lim_{L \rightarrow \infty} f_L(z) = \sum_{n=1}^{\infty} \frac{z^n}{(2\pi n)^{d/2}} e^{-n \sigma^2/2}. \quad (19)$$

This can be done using the asymptotic behavior of the E_k^L for

large L given by (8), ..., (11). Since it follows from (1) and (2) that for each $L > 0$ and $\rho > 0$ there is always a unique $\xi(L) \in [0, 1]$ we find that if $\xi(L)$ tends to $\xi \in [0, 1]$ for $L \rightarrow \infty$, then

$$\rho = \sum_{n=1}^{\infty} \frac{\xi^n}{(2\pi n)^{d/2}} e^{-n \sigma^2/2} + \lim_{L \rightarrow \infty} \frac{1}{L^d} \sum_{n=1}^{\infty} \xi^n \sum_{\{k: k_1=1\}} e^{-n \eta_k^L}. \quad (20)$$

Since

$$\lim_{L \rightarrow \infty} \frac{1}{L^d} \sum_{n=1}^{\infty} \xi^n \left[\sum_{k=1}^{\infty} \exp \left(-\frac{n \pi^2}{2L^2} (k^2 - 1) \right) \right]^{d-1} < \lim_{L \rightarrow \infty} \frac{1}{L^d} \sum_{n=1}^{\infty} \xi^n \left(1 + \frac{L}{(2\pi n)^{1/2}} \right)^{d-1} = 0,$$

We arrive at Eq. (12). Suppose now that $\xi(L) \rightarrow 1$; then

$$\frac{1}{L^d} \sum_{n=1}^{\infty} [\xi(L)]^n \sum_{\{k: k_1=1\}} e^{-n \eta_k^L} \rightarrow \rho - \rho_c. \quad (21)$$

Equation (21) implies that for $d = 1$ we have macroscopic occupation of the ground state. So for $d = 2$ we have

$$\frac{1}{L^d} \sum_{k=1}^{\infty} \frac{\xi(L)}{e^{\pi^2(k^2-1)/2} - \xi(L)} \rightarrow \rho - \rho_c, \quad (22)$$

from which it follows that $\xi(L) \sim 1 - 1/AL^2$, where A is the solution of (14). For $d \geq 3$ we have the following estimate:

$$\begin{aligned} \frac{1}{L^d} \sum_{\{k: k_1=1\}} \frac{\xi(L)}{e^{\eta_k^L} - \xi(L)} &= \frac{\xi(L)}{1 - \xi(L)} \\ &= \frac{1}{L^d} \sum_{\left\{ \substack{k: k_1=1 \\ k \neq (1, \dots, 1) \end{array} \right\}} \frac{\xi(L)}{e^{\eta_k^L} - \xi(L)} \\ &\leq \frac{1}{L^d} \sum_{\left\{ \substack{k: k_1=1 \\ k \neq (1, \dots, 1) \end{array} \right\}} \frac{1}{e^{\eta_k^L} - 1} \\ &\leq \frac{d}{L^d} \sum_{n=1}^{\infty} \sum_{k=2}^{\infty} \exp \left(-\frac{n \pi^2}{2L^2} (k^2 - 1) \right) \\ &\quad \times \left[\sum_{k=1}^{\infty} \exp \left(-\frac{n \pi^2}{2L^2} (k^2 - 1) \right) \right]^{d-2} \\ &\leq \frac{d}{L^d} \sum_{n=1}^{\infty} e^{-n \pi^2/L^2} \frac{L}{(2\pi n)^{1/2}} \left(1 + \frac{L}{(2\pi n)^{1/2}} \right)^{d-2} \\ &\leq \frac{d 2^d}{L^d} \sum_{n=1}^{\infty} e^{-n \pi^2/L^2} \frac{L}{(2\pi n)^{1/2}} \left(1 + \frac{L^{d-2}}{(2\pi n)^{d-2/2}} \right), \end{aligned} \quad (23)$$

which goes to zero as $L \rightarrow \infty$. So for $\rho > \rho_c$

$$\frac{1}{L^d} \frac{\xi(L)}{1 - \xi(L)} \rightarrow \rho - \rho_c, \quad (24)$$

and we have macroscopic occupation of the ground state only (condensation of type I).

ACKNOWLEDGMENTS

The author wishes to thank Professor Dr. J. T. Lewis for his helpful discussions. The work in this paper is part of the research program of the Foundation of Fundamental Research on Matter, which is financially supported by the Netherlands Organization for Pure Research.

- ¹M. van den Berg and J. T. Lewis, "On the free boson gas in a weak external potential," *Commun. Math. Phys.* **81**, 475 (1981).
²M. van den Berg and J. T. Lewis, "On generalized condensation in the free boson gas," *Physica A* (to be published).
³R. M. Ziff, G. M. Uhlenbeck, and M. Kac, *Phys. Lett.* **32**, 171 (1977).

- ⁴D. W. Robinson, *Commun. Math. Phys.* **50**, 53 (1976).
⁵L. J. Landau and I. F. Wilde, *Commun. Math. Phys.* **70**, 43 (1979).
⁶A. Bijl, J. de Boer, and A. Michels, *Physica* **8**, 655 (1941).
⁷J. T. Lewis and J. V. Pulè, *Commun. Math. Phys.* **36**, 1 (1974).